

Index

- Adenovirus, 137–141
Amantadine hydrochloride, 53–55, 344–345
Antigenic drift, 157–159, 194–203, 320–322, 327–334
Antigenic shift, 156–157, 320, 325–327
Antitermination during template RNA synthesis, 23, 98–106
- Bicistronic NA/NB messenger RNA (mRNA), 33–35, 38–39
Brefeldin A, 408
- Cap-dependent endonuclease, 91–92
Capped host cell primers in viral mRNA synthesis, 89, 91–92
Cleavage activation of hemagglutinin, 12, 19, 154–155
Cytotoxic T lymphocytes, 26–27, 365, 391–415
 comparison with helper T lymphocytes, 365, 412–414
 effect of major histocompatibility complex (MHC), 364, 408–410
 form of viral antigen recognized, 399–408
 recognition of individual viral proteins, 26–27, 391–399
 role in antiviral immunity, 410–412
- Defective-interfering (DI) particles, 269–311; *see also* Defective-interfering (DI) RNAs
 possible role in pathogenicity, 304–308
 possible role in virus evolution, 308–310
Defective-interfering (DI) RNAs, 269–311
 evolution, 285–288
 generation, 281–285
- Defective-interfering (DI) RNAs (*cont.*)
 mechanism of interference, 297–285
 replication, transcription and translation, 288–297
 structure, 272–281
Double-stranded RNA activated kinase (P68)
 suppression of activation in infected cells, 139–142
- Endoplasmic reticulum, 18, 38, 50, 225–242
Enterovirus 70 (EV 70), 349
Epidemiology, 319–322
Evolution of virus genes
 comparison of influenza A, B, and C viruses, 340, 349–350
 influenza A viruses, 327–336
 influenza B viruses, 338–339
 influenza C viruses, 336–338
 possible role of defective-interfering particles, 308–310
Exocytic pathway of viral glycoprotein, 18–19, 222–253
 insertion into plasma membrane, 252–254
 involvement in antigen processing, 407–408
 polarized transport, 248–252
 trans-reticular network, 19, 248–252
 translocation across the membrane of the endoplasmic reticulum, 18, 38, 50, 225–231
 transport and processing within the Golgi apparatus, 18–19, 243–248
 transport between the endoplasmic reticulum and the Golgi apparatus, 18, 241–242

- Gene mapping, 2–6
- Glycosylation of viral glycoproteins, 17, 18–19, 30, 155, 185–186, 226, 230–231, 233–235, 243–248
- Golgi apparatus, 18–19, 241–248
- Helper T lymphocytes, 365–391, 412–415
 - comparison with cytotoxic T lymphocytes, 365, 412–414
 - effect of major histocompatibility complex (MHC), 382–383
 - recognition of individual viral proteins, 375–376
 - requirement for viral antigen processing, 383–388
 - role in antiviral immunity, 388–391
- Hemagglutinin (HA), 11–20, 153–169, 219–255
 - antigenic variation, 17–18, 153–159
 - cleavage activation, 19, 154–155
 - conformational change, 20, 165–168
 - fusion properties, 20, 164–168
 - gene structure, 12–16
 - gene variation, 319–323, 325–330
 - receptor binding, 159–164
 - recognition by cytotoxic T lymphocytes, 394–395
 - recognition by helper T lymphocytes, 376–382
 - role in pathogenicity, 342–343
 - synthesis and intracellular transport, 18–19, 219–255
 - three-dimensional structure, 16–18, 153–156
- Hemagglutinin-esterase (HE), influenza C virus, 2, 20–21, 176–177, 322–323
- Influenza A viruses: *see under specific topics*
- Influenza B viruses
 - epidemiology, 320–321
 - gene variations and evolution, 338–340
 - hemagglutinin, 20, 322–323
 - membrane (M1) protein, 56–57, 323–324
 - mRNA splicing, 65–66, 131–133
 - neuraminidase, 179–180
 - nonstructural (NS1 and NS2) proteins, 65–66, 131–132, 324–325
 - nucleocapsid protein, 27, 323
 - polymerase proteins, 10
- Influenza C viruses
 - epidemiology, 320–322
 - gene variations and evolution, 336–338, 340
- Influenza C viruses (*cont.*)
 - hemagglutinin-esterase (HE), 20–22, 176–177, 181–182, 322–323
 - membrane (M1) protein, 57–58
 - mRNA splicing, 57–58, 131–133, 324
 - nonstructural (NS1 and NS2) protein, 66–67, 324–325
 - nucleocapsid protein, 27–28, 323
 - polymerase proteins, 10–11
- Interferon
 - induction of P68 kinase, 140
 - induction of the Mx protein, 116–120, 122
 - role in protective effects of cytotoxic T lymphocytes, 365, 411–412
- Internal deletions in defective-interfering (DI) RNAs, 274–285
- Jumping polymerase model, 282–284
- M2 protein, 43–56
 - gene variations, 55–56, 323–324, 334
 - function, 52–55, 344–345
 - mRNA generated by splicing, 39–42, 90, 123–124, 131
 - recognition by cytotoxic T lymphocytes, 396–397
 - structure and subcellular localization, 43–52
- Major histocompatibility complex (MHC), 364, 382–383, 408–410
- Maximum parsimony procedure, 328–338
- Membrane (M1) protein
 - functions, 42–43, 115–116
 - gene structure, 39–42
 - gene variations, 55–56, 323–324, 334
 - influenza B virus, 56–57, 132
 - influenza C virus, 57–58, 132–133
 - recognition by cytotoxic T lymphocytes, 396–397
 - role in pathogenicity, 344–345
 - time course of synthesis, 110–116
- Membrane fusion by the hemagglutinin, 20, 164–165
- Methylation of internal adenosine (A) residues of viral mRNA, 90, 123
- Mutation rates for RNA viruses, 345–347
- Mx protein, 116–123
- NB protein, influenza B virus, 34–49
- Neuraminidase (NA), 28–34, 175–210, 219–255
 - antigenic sites, 194–209
 - comparison of influenza virus A and B strains, 30–31, 182–188, 330–331

- Neuraminidase (NA) (*cont.*)
 function, 28, 175–178
 gene structure, influenza A virus, 29–30, 179–180
 gene structure, influenza B virus, 32–34, 179–180
 membrane association, 28–29, 31–32, 178–179
 role in pathogenicity, 343–344
 synthesis and intracellular transport, 219–255
 three-dimensional structure, 32, 188–194
 three-dimensional structure of complexes with antibody, 205–209
- Nonstructural protein 1 (NS1)
 function, 62, 109–110
 gene structure, 62, 58–59, 62–63, 123
 influenza B virus, 65–66, 131–132
 influenza C virus, 66–67
 nuclear transport sequences, 63–64
 recognition by cytotoxic T lymphocytes, 397
 time course of synthesis, 110–116
- Nonstructural protein 2 (NS2)
 influenza B virus, 65–66, 131–132
 influenza C virus, 66–67
 mRNA generated by splicing, 58–61, 123–131
 recognition by cytotoxic T lymphocytes, 397
 structure, subcellular localization and function, 65–66, 109–110
- Nuclear transport (karyophilic) sequences, 9, 24–25, 63–64, 122
- Nuclear-cytoplasmic transport of virus-specific RNAs, 114–115
- Nucleocapsid (NP) protein
 association with viral nucleocapsids (or ribonucleoprotein particles), 22–23, 91–92
 gene variations, 323, 333
 influenza B virus, 27
 influenza C virus, 27–28
 nuclear transport sequences, 24–25
 recognition by cytotoxic T lymphocytes, 26–27, 396
 roles in viral RNA replication, 101–110, 115–116
 structure, 23–24
 time course of synthesis, 110–116
- Paramyxovirus HN protein, 31, 181
- Pathogenicity, 341–345
- Pathogenicity (*cont.*)
 possible role of defective-interfering particles, 304–308
- Polarized transport of viral glycoproteins, 248–252
- Poliiovirus, 346–348
- Polymerase (P) proteins, 6–11, 91–98; *see also* Viral RNA transcriptase
 formation of complexes, 8–10, 93–96
 gene deletions in defective-interfering RNAs, 269–281
 gene structure, 608
 influenza B virus, 10
 influenza C virus, 10–11
 internally deleted forms encoded by defective-interfering (DI) RNAs, 289–297
 recognition by cytotoxic T lymphocytes, 396
 roles in mRNA synthesis, 8, 91–93
 roles in replication, 108
 structure of internally deleted genes in defective-interfering (DI) particles, 274–281
- Polysomes in infected cells, 135–139
- Positive selection of virus variants, 335, 349–350
- RNA polymerase II transcripts
 metabolism in infected cells, 133–134
 source of capped RNA primers, 89, 91–92
- Receptor binding, 159–164
- Retroviruses, 90, 131, 346–348
- Reye's syndrome, 320–321
- Rollover/loop-out polymerase model, 282–284
- Shutoff of host cell translation, 133–139
- Sialic acid receptors, 11, 159–164
- Signal recognition particle (SRP), 38, 50–51, 225
- Signal sequence of hemagglutinin, 12, 18, 225–226
- Small nuclear ribonucleoproteins (snRNPs), 125, 126–130
- Spliceosomes, 125, 126–130
- Splicing of viral messenger RNAs (mRNAs)
in vitro studies, 124–131
 NS1 mRNA of influenza A virus, 58–61, 123–131
 M1 mRNA of influenza A virus, 39–42, 90, 123–124, 131
 NS1 mRNA of influenza B virus, 65–66, 131–132

- Splicing of viral messenger RNAs (*cont.*)
 M1 mRNA of influenza B virus, 56–57, 132
 M1 mRNA of influenza C virus, 131–133, 324
- Temperature-sensitive virus mutants
 hemagglutinin, 236–237, 251
 neuraminidase, 236–237
 nucleocapsid protein, 103–105
 polymerase, 9, 92, 93
- Template RNA synthesis
 mechanism, 98–106, 108, 109–110
 regulation in infected cells, 110–116
Trans-reticular network, 19, 248–252
- Translation initiation factors eIF-2 and eIF-2B, 139–142
- Translation initiation
 bicistronic NA/NB mRNA, 33–35, 38–39
 in influenza virus-infected cells, 135–142
- Translocation across the membrane of the endoplasmic reticulum, 18, 38, 50, 225–231
- Variations in nature of human RNA viruses, 347–349
- Vesicular stomatitis virus (VSV), 23, 97, 116
- Viral RNA transcriptase, 22–23, 91–98
- Viral messenger RNA (mRNA) synthesis
 inhibition by the murine Mx protein, 120–122
 mechanism, 89, 91–98
 regulation in infected cells, 110–116
- Viral nucleocapsids
 assembly, 25–26, 93–98, 104, 108–109
 function, 23, 91–92, 108
 structure, 22, 91–93
see also Viral RNA transcriptase; Viral ribonucleoproteins (RNPs)
- Viral protein synthesis
 regulation in infected cells, 110–116
 selective and efficient translation of viral mRNAs, 133–139
 suppression of double-stranded RNA-activated kinase (P68), 139–142
- Viral ribonucleoproteins (RNPs): *see* Viral nucleocapsids
- Virion assembly, 26, 43, 253–254
- Virion RNA synthesis
 mechanism, 106–110
 regulation in infected cells, 110–116
- Virulence, 341–345